

IPB Seminar Series in Plant Biochemistry

Cell-to-cell transport in differentiated roots

Dr. Marie Barberon
University of Geneva, Switzerland

Thursday 08th of February, 2024, at 4 p.m.

Kurt Mothes Hall, Leibniz Institute of Plant Biochemistry

Plant roots forage the soil to acquire water and nutrients for growth and development. This function is closely linked to their anatomy: water and nutrients move radially through the concentric layers of epidermis, cortex, and endodermis before entering the vasculature. This arrangement allows for three uptake scenarios: the “symplastic pathway”, where the outer cells actively take up nutrients, which are then transported from cell to cell through plasmodesmata; the “apoplastic pathway”, where nutrients are transported in the apoplast and blocked by the endodermal apoplastic barrier (Caspasian strips); and the “coupled trans-cellular pathway”, where nutrients are transported sequentially from one cell to another by polarized influx and efflux carriers and are barred by the endodermal diffusion barrier (suberin lamellae). My group aims to functionally characterize these pathways for nutrient acquisition by a combination of physiology, cell biology and developmental approaches. We are particularly interested in suberin lamellae and plasmodesmata function and regulations for nutrient acquisition. Our recent work identified key regulators of suberin deposition and cell-to-cell transport in roots.

Host:

Dr. Christin Naumann
Nutrient Sensing
IPB, Halle (Saale)



Leibniz-Institut für
Pflanzenbiochemie

cnaumann@ipb-halle.de

Tel. +49 345 5582 1220

